



Quality of M-5 Agronomy Characteristics of Black Rice Induction Mutation to Get a Short Rods

Benny Warman¹; Hendra Alfi¹; Veronice²

¹ Lecturer of Food Security Studi Program, Politeknik Pertanian Negeri Payakumbuh, Indonesia

² Lecturer of Agribusiness Management Studi Program, Politeknik Pertanian Negeri Payakumbuh, Indonesia

<http://dx.doi.org/10.18415/ijmmu.v10i1.4256>

Abstract

Local black rice is a genetic potential that is very potential and strategic to be developed in the future. In fact, if the short comings (production age > 165 days and higher > 150 cm) can be improved it will obviously be better, both in economic value and in terms of supporting the national food security program. Therefore, the effort to preserve and genetic improvement of local black rice which is a genetic wealth asset needs to be done. The purpose of this study was to obtain an overview of agronomic characters and production potential in several semi-dwarf lines of black rice. In this study, through gamma-induced mutation 18 strategic mutant rice lines have been produced with strategic expectations (more early maturing / 30-40 days faster harvest compared to the original variety, and lower / 50-65 cm posture) which are the result of improvement. genetic local black rice from West Sumatra. These strategic expectations of mutant rice strains have pera taste preferences, so that they have the taste preferences of the people of West Sumatra.

Keywords: *Black Rice; Mutation; Short Rods; West Sumatera Landrace; Induced Mutations; Early Maturity*

Introduction

The use of improved varieties is a reliable technology in improving agricultural production. This technology is seen as safer and more environmentally friendly and less costly for farmers. But in generating improved varieties need to be carried out a series of breeding activities so that the expected benefits can be generated. Until now, various activities to produce rice varieties continue to be made. However, most of the improved varieties that produced less in demand by the people of West Sumatra because mismatches taste preferences. This causes people in West Sumatra are still a lot of local rice cultivars cultivated than modern varieties

Recently in West Sumatra there are more than 50 local rice (landraces) cultivars, both paddy and upland rice [5], and some of them are already known by the public and has a pretty good economic value. Generally, the character of the local rice cultivars in the old (≥ 120 days). When the age of the local rice

can be shortened (early maturing) would be better, in particular can increase the intensity as well as the annual crop can be more profitable for farmers.

Efforts for improvement Local rice genetic should be made to produce superior crops including early maturity. One way that can be done in the development of improved varieties is by mutation induced by gamma ray irradiation. Mutation breeding can support the resulting large genetic variability of plants that have a narrow genetic variability [20], [19], [10]. In addition, through mutation breeding can also be generated large genetic variability within a shorter time when compared with hybridization methods.

Mutation induction is one way to change the genetic done by humans in order to obtain better properties than the original plant properties [10], [26], [13]. Mutation induction has a significant contribution to the genetic improvement of plants in various parts of the world. In fact, at some point have an impact on the increase in production as well as rice [18], it is better to repair some properties just by not changing the majority of plant traits original that has been favored and relatively requires less time in the refining process strains [19], [2], [14]. The research objective is to get as a result of early maturing mutant local rice genetic improvement through mutation breeding of West Sumatra.

Materials and Methods

This research was conducted in the greenhouse facilities and rice experiment Polytechnic State Agriculture Payakumbuh since April 2009 to December 2010. The plant material used is Junjung varieties which is one of the local rice cultivars West Sumatra. Seed blue label irradiated with a beam of γ with doses of 0 Gy (control), 100 Gy, 200 Gy, 300 Gy, 400 Gy, 500 Gy, 600 Gy, 700 Gy, 800 Gy, 900 Gy, and 1000 Gy of 100 grams per dose as seed M_1 . Seed irradiation is done in PAIR-BATAN, Pasar Jumat Jakarta with γ -ray radiation source (gamma) used Co^{60} of Gammacell Chamber. Once irradiated, each irradiation dose of 200 seeds planted in the seedbed for three weeks to observe the effect of irradiation on the growth pattern of the seeds and determine the value of LD_{50} . In addition, it also made observations of chlorophyll mutations according to the **Gustafssons** method [8], to determine the frequency of mutant and mutation frequency.

Treatment dosage is best then the garden to get the source of planting material and plant strains formation at the stage of M_2 . Furthermore, each strain were planted about 100 plants per planting hole. Plants grown in paddy rice cultivation as well maintained as usual. In the generative growth phase carried out observations of genetic diversity, heritability and the selection of the target mutant (mutant-old early maturing). The determination of the genetic variability was performed according to methods Khan [16], in which a character has a genetic variability that is broad if the value of genetic variance (σ^2g) more than twice the standard deviation of genetic or expressed by the formula $\sigma^2g > 2 (\sigma\sigma^2g)$, reverse genetic variability narrow if $\sigma^2g < 2 (\sigma\sigma^2g)$. Determination of heritability done using analysis of variance components and is calculated based on the formula proposed [29] and [6] in which the heritability estimates higher when ($H > 50\%$), moderate ($20\% \leq H \leq 50\%$), and low ($H < 20\%$).

Results and Discussion

At this stage of the M_1 , from research conducted during germination until age 21 day after planted can be seen clearly seedling growth response (percentage of germination, seedling height, root length and percentage of germination) local rice cultivars West Sumatra to the irradiation dose being treated. Seedling growth response to treatment irradiation dose can be seen in Figure 1. In Figure 1a. it appears that the higher the dose of irradiation, can reduce the ability to grow crops. In fact, at doses above

500 Gy, causing death in sprouts (lethal effects). The declining vitality of the plant due to the deterministic effects due to gamma-ray irradiation.




Figure 1. The response curve gamma irradiation treatment at various doses on germination and seedling growth of cultivars Junjung. (a) the effect of irradiation on germination, (b) the effect of irradiation on seedling height, (c) the effect of irradiation on root growth. (d) the effect of irradiation on sterility seeds.

Deterministic effect is the effect caused by cell death due to radiation exposure. The deterministic effects occur when the dose received by the plant above the threshold dose (threshold dose) and generally arise some time after irradiation. Heightened reduction of growth due to the increasing size of the dose proves that the irradiation dose is very direct effect on physical damage (physical damage) plants. Damage to the M1 generation is due to ionizing radiation treatment [7]. In general, the physiological damage are more descriptive. It can be observed directly morphologically. Increased doses of mutagens causing mortality rate higher plants with a low mutation frequency. It is the result of damage extrachromosomal. Conversely if these mutagens cause mortality rates were low and the high frequency of mutation is believed to be due to chromosomal damage.

The germination percentage of observations obtained by the mathematical equation where $y = -0.135x + 117.1$, so the LD50 values obtained at 497.04 Gy irradiation dose. High-low levels of oxygen and water molecules (H_2O) in the irradiated material, will affect the process of ionization of free radicals are formed that affect radiosensivitas plant cell [11]. In addition, the lethal activity of ionizing radiation on cells resulting in the loss of mitotic activity, resulting in a loss of genetic material [1]. This leads to a low rate of germination and seedling growth and can even cause death in plant material (lethal) as shown in Figure 1a, 1b, and 1c.

At the seedling height and root length also seen that the higher the dose of irradiation also affect the growth response. High growth response of seedlings and root length decreased due to the growing size of the irradiation dose. At doses above 500 Gy just a little seed that grows and then death. This is in line with the opinion [17], [4], which explains that the treatment of certain mutagens in serealea has a high correlation with sprouts M_1 , M_1 germination and mutation frequency.

One thing that is expected in the physiological damage induced mutation is the smallest and genetic damage as much as possible. This is a very valuable factor in generating high genetic variability.

In this study the LD₅₀ values obtained are within the range of 400-500 Gy dose, but in a dose range is still causing panicle sterility levels are very high (Figure 1.d.). High doses of radiation will increase sterility in plants panicles M₁ [7]. From Figure 1d. also can be seen that for a dose of 200-300 Gy panicle sterility levels ranging between 20-30%, while the percentage of sterility dose of 0-100 Gy panicle hardly much different.

Gamma irradiation at a dose of 200 Gy can cause chlorophyll mutations in M₂ population as shown in Table 1. While the origin of the plant cultivars (control) there is no mutation of chlorophyll. According to [10], the chlorophyll mutation is an indication of genetic damage, which is characterized by the formation of chlorophyll in leaves. It was observed that at a dose of 200 Gy irradiation has produced a number of chlorophyll mutations are quite spacious with 7 different types of mutation, namely Albina, albovidis, chlorina, virescen, tigrina, marginata, and striata. Some types of mutations that appear on M₂ nurseries can be seen in Figure 2.

Frequency of chlorophyll mutants in M₂ population can also be seen in Table 1. Of the seven types of chlorophyll mutations are formed, the number of mutations towards albino higher when compared with other types of chlorophyll mutations. Chlorophyll mutation frequency obtained 0.17% and 1.10% mutant frequency. It is lower when compared to the results reported by [31] by 7.7% on the variety Hitomebore [26] of 11.2% and 6.4% on the cultivar Kuriak Kusuik and Randah Putiah. Tinggi. Chlorophyll mutations that occur in a population of M₂ derived from gamma-ray irradiation at a dose of 200 Gy. a mutagenic effect of gamma irradiation at a dose indicating that the genetic diversity of the population has created M₂. [31] and [26] also reported that a dose of 200 Gy is the irradiation dose that is effective in generating useful genetic diversity in breeding programs.



Figure 2. Movements of chlorophyll which appeared on the M₂ generation

At this stage M₂ mutant selection activity against the target (early maturity mutant) done. The selection can be done if there is a mutant strain in the population genetic diversity support to do the selection of the target mutant character [10]. Summary genetic parameter data observed through the

variable plant height, panicle length, number of productive tillers and flowering dates can be seen in Table 2. In Table 2 it appears that in general it appears that heritability estimates on the overall character including the high category. Only the character of panicle length, which is estimated heritability was moderate, but can be classified nearing high. The high heritability estimates this means genetic factors make important contributions in the next process. Heritability shows how the proportion of a gene can be passed down to the next generation is based on observation of the nature of the observed phenotype [24], [30], as well as the genetic parameters determining the selection of effective systems [25], [30].

Table 1. Mutation Type of Chlorophyll mutation, Total Mutation, Mutation Frequency and Frequency of Mutants in irradiation dose of 200 Gy

Kult. Junjung	Chlorophyll Mutation Type							Frek. Mutants (%)	Frek Mutations (%)
	Alb	Albo	Chl	Vir	Tig	Mar	Stri		
Total	993	78	567	136	6	193	3	1.10	0.17
Freq . of Mutants (%)	0.55	0.04	0.32	0.08	0.00	0.11	0.00	-	-

Description: The Alb (Albina), Albo (Alboviridis), Chl (Chlorina), Vir (virescens), Tig (Tigrina), Mark (Marginata), Stri (Striata)

Table 2. General summary of genetic parameters on the part of the M₂ population

Character	μ	Sd	$\sigma^2 P$	$\sigma^2 e$	$\sigma^2 G$	h^2	Category	2.Sd	Variability
Plant Height (cm)	84.76 [*]	±4.54	21.27	9.00	12.27	0.52	height	9.09	Broad
Tassel Height (cm)	26.48 ^{ns}	±1.69	2.90	1.46	1.44	0.46	Medium	3.37	Limited
Numbers of Productive tillers	13.63 ^{ns}	±2.37	5.71	2.13	3.58	0.60	height	4.74	Limited
Age of Flowering (Day)	94.85 ^{ns}	±2.78	10.45	1.47	9.41	0.96	height	5.55	Broad

Note : *) significantly different at the 0.05 level according to the T test;
^{ns}) were not significantly different

In addition, the value of variability in plant height and days to flowering is broad while the character panicle length and the number of productive tillers relatively narrow. This means that the induced mutation made more effective in generating high genetic diversity in plant height and days to flowering. But for the panicle length and the number of productive tillers do not have a great influence in generating genetic diversity. The extent of genetic variability in the character of the age of flowering and plant height are the basis for selection against a target mutant character (superior) which is desirable [10], [23], such as early maturing mutants and semi-dwarf. In observation of the middle value of a variable, it is known that the value of the variable character of mutant plants posture is lower when compared with the posture of plant origin. As for the value of the middle third of other variables that are not significantly different from the value of the middle of the plant origin. According to [30], the midpoint of each genotype is associated with idio-type plants to be achieved and desired as semi-dwarf character and early maturity.

In the generative phase M₂ populations do individual selection against a target mutant (mutant early maturing and semi-dwarf). Results of individual selection against mutants early ripening and semi-

dwarf can be seen in Table 3 and Table 4. In Table 3 shows that the result of individual selection of the candidate mutant very early-maturing group (age 70-80 days flowering) that is as much as 8 plants, and the group of candidates mutant early maturing (flowering age 81-90 days) as many as 78 plants, the mutant frequency of 0.14%. As for the origin of plants (control) contained in the flowering age groups of 91-100 days. Mutant very early maturing and early maturing mutants with semi-dwarf can be seen in Figure 3.




Figure 3. Mutant very early maturing and early maturing mutants with semi-dwarf stature genetic improved results through induced mutation were found in M₂ population

At doses of gamma irradiation at a dose of 200 Gy can bring out the mutant as a result of genetic changes due to mutations in genes that control the age of the plant. From observations made rapid harvest age can be characterized by more rapid flowering mutant plant age when compared with native plants (plant control). According [12] days to flowering faster highly correlated with crop harvesting age faster. The rapid flowering plants due to genetic changes as a result of mutation induction is done. Treatment irradiation dose of 200 Gy is the dose that is effective for generating mutants is strategically desirable as mutant semi-dwarf and mutant early maturing [15], mutant grain hairless varieties Cisantana [22], and Mutant early maturing on rice Kuriak Kusuik and High White more low [26], early maturing mutants and semi-dwarf in rice Ase Field and Mandoti [9].

Table 3. Early maturing mutant in M₂ population derived from cultivars Junjung

Tread	Age for Flowering (day)			Numbers of Mutans	Numbers of Populations M ₂	Frek.Early-maturing Mutans (%)
	70-80	81-90	91-100			
Mutans of Junjung	8	78	-	86	60.000	0.14
Kult. Junjung (Control)	-	-	TF 50	-	6.000	-

Note: TF50 (*Time Flowering* for Plant Control 50%)

In addition to producing the mutant-old early maturing, irradiation necessary also produce mutants dwarf and semi-dwarf. Results of selection of mutant dwarf and semi-dwarf can be seen in Table 4. Generally, in early maturing mutant phenotype candidate has a lower posture (semi-dwarf) when compared with native plants idiotipe. When connected with genetic parameters, in addition to the character of the age of flowering plants, plant height also has a wide genetic diversity value (see Table 2).

Table 4. Mutant dwarf and semi-dwarf in M₂ population of derived from cultivars Junjung

Thread	Postur of Plant		Numbers of Mutans	Numbers of Populations M ₂	Frek.Early-ripening Mutans (%)
	<i>Dwarf</i>	<i>Semi-Dwarf</i>			
Mutans of Junjung	27	70	97	60.000	0.16
Frek. Mutants (%)	0.05	0.12			

In Table 4 it is known that the mutant dwarf mutants obtained as many as eight candidates and semi-dwarf mutants obtained as many as 70 candidates mutant with a mutation frequency of 0.16%. This frequency is lower than the frequency of mutant dwarf and semi-dwarf those reported by [3] in the amount of 0.30% on the cultivar Madhu Malti at a dose of 250-350 Gy irradiation, and [28], which is 0.26% in strain KI 237 with irradiation dose of 200 Gy.

Conclusion

Based on the research that has been done can be concluded that the mutation induction is done by gamma ray irradiation can generate genetic diversity is very useful in the selection efforts to obtain the characters age more early maturing crops (early maturing mutant). Orientation doses of LD₅₀ values obtained at 497.04 Gy, but in determining the expected effective dose irradiation doses causing little physical damage as well as a large genetic damage. At a dose of 200 Gy irradiation produced little physical damage as a percentage of germination, seedling height, root length and panicle void percentage when compared with the higher dose of irradiation and generate genetic diversity seen from the emergence of chlorophyll mutation with a mutation frequency of 0.17%. At these doses also has established extensive genetic diversity for the variable character of the age and height of the plant. With the extent of genetic diversity of the resulting mutant candidates that can be selected at this stage of M₂. From the selection made at the stage of early maturing mutant M₂ candidates obtained as many as 86 candidates with the mutation frequency of 0.14%, as well as dwarf mutant and semi-dwarf mutant were 97 mutants with mutant frequency of 0.16%, Candidates mutant is a mutant candidates hope to have better character as a result to be developed in the future.

References

- [1] Ahnstrom, G. 1977. Radiobiology. In Manual on Mutation Breeding. 2nd Ed. IAEA, Vienna. 21-27.
- [2] Amano, E. 2004. Practical suggestion for mutation breeding. Forum for Nuclear Cooperation in Asia (FNCA) Mutation Breeding Project.
- [3] Bansal, V.K., and P.C. Katoch. 1991. Selection of semi-dwarf, early maturing and blast resistant mutants after mutagenic seed treatment in two locally adapted Indian Rice cultivars. Plant Breeding 107 (169).
- [4] Conger, B.V., Konzak, C.F. and R.A. Nilan. 1977. Radiation sensitivity and modifying factors. In Manual on AMutation Breeding, 2nd Ed. IAEA, Vienna. 40-42.
- [5] Etti Swasti, I. Suliansyah dan A.A. Syarif. 2007. Eksplorasi, identifikasi dan pemantapan koleksi plasmanutfeh padi asal Sumatera Barat. Lembaga Penelitian Universitas Andalas Padang.

- [6] Falconer, D.S. and T.F.C. Mackay. 1996. Introduction to Quantitative Genetics. 4th ed. Addison Wesley Longman. Harlow, Essex. UK.
- [7] Gaul, H. 1977. Mutagen effects in the first generation after seed treatment. In Manual on Mutation Breeding. 2nd Ed. IAEA, Vienna. 40-42.
- [8] Gustafsson, Å. 1938. Studies on the genetic basis of chlorophyll formation and the mechanism of induced mutagenesis. *Hereditas*, 24: 33-93.
- [9] Haris, A., Abdullah, Bakhtiar, Subaedah, Aminah and K. Jusoff. 2013. Gamma ray radiation mutant rice on local dwarf. *Middle-East Journal of Scientific Research* 15 (8).
- [10] Harten, A.M.van. 1998. Mutation breeding; Theory and practical application. Cambridge University Press.
- [11] Herison, C., Rustikawati, Sujono H. S., Syarifah I. A. 2008. Induksi mutasi melalui sinar gamma terhadap benih untuk meningkatkan keragaman populasi dasar jagung (*Zea mays* L.). *Akta Agrosia* 11(1):57-62.
- [12] Ismachin, M. 1983. Sifat genjah mutan padi varietas Pelita I/1 dan IR5. Disertasi Doktor. Disampaikan pada Fakultas Pascasarjana IPB. Bogor (Tidak Dipublikasikan).
- [13] Ismachin, M. 2007. Perkembangan pemuliaan mutasi di Indonesia. Diklat Pemuliaan Mutasi. FPAI BATAN. Jakarta.
- [14] Ismachin, M. dan Sobrizal. 2006. A significant contribution of mutation techniques to rice breeding in Indonesia. *Plant Mutation Report* Vol. 1, No. 18.
- [15] Juliano, B.O. and C.P. Villareal. 1983. Grain quality evaluation of word rices. IRRI, Los Banos, Manila. Phillippines.
- [16] Khan, A.S., M. Imran, M. Ashfaq. 2009. Estimation of genetic variability and correlation for grain yield component in rice (*Oryza sativa* L.). *Am-Euras. J. Agric. Environ. Sci.* 6:585-590.
- [17] Konzak, C.F., R.A. Nilan., J. Wagner., and R.J. Foster. 1965. Efficient chemical mutagenesis Report of FAO/IAEA Technical Meeting. Rome, May 25 – June 1st 1964. 49-70.
- [18] Maluszinsky, M., B.S. Ahloowalia, B. Sigurbjornsson. 1995. Application of in vivo and in vitro mutation techniques for crop improvement. *Euphytica* Vol. 85 (303).
- [19] Micke, A., B. Donini, M. Maluszinski. 1990. Induced mutation for crop improvement. *Mutation Breeding Review* Vol. 7. No. 1.
- [20] Mikaelson, k. 1980. Modifying effects of water content, oxygen and cystein on genetics effects of gamma and fast neutron radiation in barley seeds. In *Gamma Field Symposia*: 7:1-16.
- [21] Micke, A., B. Donini, M. Maluszinski. 1990. Induced mutation for crop improvement. *Mutation Breeding Review* Vol. 7. No. 1.
- [22] Mugiono, L. Harsanti, dan A.K. Dewi. 2009. Perbaikan padi varietas Cisantana dengan mutasi induksi. *Jurnal Ilmiah Aplikasi Isotop dan Radiasi* Vol. 5 (2.).

- [23] Roy, D. 2000. *Plant Breeding: Analysis and Exploitation of Variation*. Narosa Publishing House Calcutta.
- [24] Sabu, K.K., M.Z. Abdullah, L.S. Lim, R. Wickneswari. 2009. Analysis of heritability and genetic variability of agronomically important traits in *Oryza sativa* L. x *O. rufipogon* Cross. *Agronomy Res.* 7:97-102.
- [25] Susanto, G.W.A. dan M.M. Adie. 2010. Pendugaan heritabilitas hasil dan komponen hasil galur-galur kedelai ditiga lingkungan. *Prosiding Simposium PERIPI 5-7 Agustus 2004.*, Hal: 119-125.
- [26] Sobrizal. 2007. Seleksi mutan genjah pada populasi M2 tanaman padi varietas Kuriak Kusuik dan Randah Tinggi Putih. *Jurnal Agrotropika* Vol.1 No.
- [27] Sobrizal dan Ismachin, M. 2006. Peluang mutasi induksi pada upaya pemecahan hambatan peningkatan produksi padi. *Jurnal Ilmiah Aplikasi Isotop dan Radiasi*. Vol. 2, No. 1. BATAN. Jakarta.
- [28] Sobrizal. 2008. Mutasi induksi untuk mereduksi tinggi tanaman padi galur KI 237. *Jurnal Ilmiah Aplikasi Isotop dan Radiasi* Vol. 4, No. 2.
- [29] Stansfield, W.D. 1983. *Theory and Problems of Genetic*. 2nd Ed. New York. McGraw-Hill.
- [30] Syukur, M., S. Sujiprihati., R. Yuniarti., dan D.A. Kusumah. Estimation of genetic variance and heritability for yield component characters in chili pepper genotypes. *J. Agrivigor*10(2): 148-156.
- [31] Yamaguchi, H., T. Morishita, K. Degi, A. Tanaka, N. Shikazono, and Y. Hase. 2006. Effect of carbon-ion beams irradiation on mutation induction in rice. *Plant Mutation Report*. 1:25-27.

Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal.

This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).